

Arteriosclerosis, Thrombosis, and Vascular Biology

JOURNAL OF THE AMERICAN HEART ASSOCIATION

American Heart
Association®



Learn and Live SM

From Vessel Sprouting to Normalization. Role of the Prolyl Hydroxylase Domain Protein/Hypoxia-Inducible Factor Oxygen-Sensing Machinery

Cathy Coulon, Maria Georgiadou, Carmen Roncal, Katrien De Bock, Tobias Langenberg and Peter Carmeliet

Arterioscler Thromb Vasc Biol published online Oct 21, 2010;

DOI: 10.1161/ATVBAHA.110.214106

Arteriosclerosis, Thrombosis, and Vascular Biology is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75214

Copyright © 2010 American Heart Association. All rights reserved. Print ISSN: 1079-5642. Online ISSN: 1524-4636

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://atvb.ahajournals.org>

Subscriptions: Information about subscribing to Arteriosclerosis, Thrombosis, and Vascular Biology is online at

<http://atvb.ahajournals.org/subscriptions/>

Permissions: Permissions & Rights Desk, Lippincott Williams & Wilkins, a division of Wolters Kluwer Health, 351 West Camden Street, Baltimore, MD 21202-2436. Phone: 410-528-4050. Fax: 410-528-8550. E-mail:

journalpermissions@lww.com

Reprints: Information about reprints can be found online at

<http://www.lww.com/reprints>

From Vessel Sprouting to Normalization

Role of the Prolyl Hydroxylase Domain Protein/Hypoxia-Inducible Factor Oxygen-Sensing Machinery

Cathy Coulon, Maria Georgiadou, Carmen Roncal, Katrien De Bock,
Tobias Langenberg, Peter Carmeliet

Abstract—The accepted model of vessel branching distinguishes several endothelial cell fates. At the forefront of a vessel sprout, “tip cells” guide the sprouting vessel toward an angiogenic stimulus. Behind the tip, “stalk cells” proliferate to elongate the vessel branch and create a lumen. In mature vessels, endothelial cells acquire a streamlined shape to optimally conduct blood flow. For this purpose, endothelial cells switch to the “phalanx” cell fate, which is characterized by quiescent and nonproliferating cells aligned in a tight cobblestonelike layer. Vessel maturation also requires the recruitment of mural cells (ie, smooth muscle cells and pericytes). These cell fates are often altered in pathological conditions, most prominently during the formation of tumor vasculature. Given the essential role of hypoxia as the driving force for initiating angiogenesis, it is not surprising that the hypoxia-sensing machinery controls key steps in physiological and pathological angiogenesis. (*Arterioscler Thromb Vasc Biol.* 2011;31:00-00.)

Key Words: angiogenesis ■ hypoxia ■ HIF ■ PHD

Angiogenesis is a dynamic process that is tightly regulated by oxygen-sensing mechanisms. The prolyl hydroxylase domain proteins (PHDs) are oxygen-sensing enzymes that belong to the evolutionarily conserved superfamily of iron-containing 2-oxoglutarate-dependent dioxygenases. This family consists of 3 PHDs (PHD1–3) and the factor inhibiting hypoxia-inducible factor (HIF) (FIH).¹ Under well-oxygenated conditions, PHDs hydroxylate conserved prolyl residues of the HIF- α subunits, thereby generating a binding site for the von Hippel–Landau tumor suppressor protein, which targets HIF- α for proteosomal degradation. When oxygen levels decrease, the hydroxylation activity of the PHDs is reduced, leading to HIF- α accumulation. On binding to HIF- β , the HIF- α/β complex translocates and activates transcription of numerous genes, including those regulating survival, metabolism, and angiogenesis.² In normoxia, the transcriptional activity of HIF is reduced by hydroxylation of a conserved asparagine residue by FIH.³ Emerging evidence suggests that HIF-1 α and HIF-2 α have distinct nonoverlapping roles in angiogenesis.⁴ PHD2 is a key family member and, consistent with this report, its deficiency disrupts development.^{5,6} Oxygen-independent pathways for the regulation of HIFs and PHDs also exist.^{1,7} Moreover, the oxygen sensors can hydroxylate other proteins, some of which are associated with angiogenesis.¹ Important knockout phenotypes of these PHD/HIF molecules in mice are listed in the Table.

Oxygen Sensing in the Endothelial Sprout

Vessel branching is regulated by growth factors^{8,9} and guidance cues.^{10,11} Gradients of some of these signals are sensed by the tip cell, which translates them into directional migration. Tip cells are polarized, have numerous filopodia that probe the environment while the cell migrates toward an angiogenic stimulus, and do not form a lumen.⁸ They have a specific molecular signature, characterized by the expression of vascular endothelial growth factor (VEGF) receptor 2 (VEGFR2), VEGFR3, platelet-derived growth factor- β , Unc5B, neuropilin, delta-like ligand-4 (Dll4), CXCR4, and Ephrin-B2,^{12–15} among other markers.¹⁶ Dll4 binds to the Notch1 receptor of neighboring cells, thereby inducing a stalk cell phenotype.¹⁷ Stalk cells trail behind the tip, proliferate, elongate the stalk, form a lumen, and connect to the circulation. They do not extend filopodia but rather form tight junctions and prevent endothelial cell (EC) retraction.¹⁶

Although hypoxia is a strong stimulus for angiogenesis, it remains unclear how hypoxia-induced signaling regulates the various steps of vessel branching. Hypoxia and HIF-1 α enhance the expression of VEGFR2,⁸ which induces Dll4 expression in the tip cell. Components of the oxygen-sensing machinery also modulate the Notch signaling pathway. For instance, HIF-1 α interacts with the Notch intracellular domain and increases its activity.¹⁸ Moreover, HIFs stimulate transcription of the Notch targets Hey1, Hey2, and Dll4,^{19,20} whereas HIF-1 α heterozygosity decreases the levels of acti-

Received on: August 1, 2010; final version accepted on: October 7, 2010.

From the Vesalius Research Center, KU Leuven, Leuven, Belgium; and the Vesalius Research Center, VIB, Leuven, Belgium.

Correspondence to Peter Carmeliet, MD, PhD, Vesalius Research Center, VIB, KU Leuven, Campus Gasthuisberg, Herestraat 49, B-3000 Leuven, Belgium. E-mail: peter.carmeliet@vib-kuleuven.be

© 2010 American Heart Association, Inc.

Arterioscler Thromb Vasc Biol is available at <http://atvb.ahajournals.org>

DOI: 10.1161/ATVBAHA.110.214106

Table. Summary of Phenotypes in Knockout Mice of PHD/HIF Oxygen-Sensing Molecules

Gene	Genotype	Phenotype
<i>PHD1</i>	−/−	Hypoxia tolerance of skeletal muscle and liver due to reprogramming to anaerobic metabolism ^{72,73} and decreased cyclin D1 levels and mammary gland proliferation ⁷⁴
<i>PHD2</i>	+/-	Normal vascular development; tumor vessel normalization, leading to reduced metastasis and tumor invasiveness due to improved tumor oxygenation ³⁰
	−/− (Prenatal)	Embryonic lethality due to placental and heart defects ⁵
	−/− (Postnatal)	Increased angiogenesis ⁷⁵ and polycythemia and congestive heart failure ⁷⁶
	Silencing in tumor cells	Increased tumor growth and angiogenesis ⁴²
	Endothelial haplodeficiency	Tumor vessel normalization (phenocopy of the <i>PHD2</i> ^{+/-} phenotype) ³⁰
<i>PHD3</i>	−/−	Hypofunctional sympathoadrenal system and reduced blood pressure ⁷⁷
<i>PHD1; PHD3</i>	−/−; −/−	Viable and fertile with smaller litters and moderate erythrocytosis ⁴³
<i>PHD3; PHD2</i>	−/−; −/− (Postnatal)	Premature lethality due to hepatic steatosis and dilated cardiomyopathy ⁷⁸
<i>FIH</i>	−/−	Viable, but reduced, body mass; enhanced metabolic rate; and insulin sensitivity ⁷⁹
<i>HIF-1α</i>	+/-	Protection against vascular remodeling during pulmonary hypertension and impaired physiological response to chronic hypoxia ⁸⁰
	Endothelial inactivation	Impaired tumor angiogenesis ²⁵
	<i>HIF-1α</i> ^{KI/+} : <i>HIF-2α</i> knocked in <i>HIF-1α</i> allele	Viable and decreased incidence of thymic lymphomas in a p53 mutant mouse model ²¹
	<i>HIF1α</i> ^{KI/KI} : <i>HIF-2α</i> knocked into both <i>HIF-1α</i> alleles	Embryonic lethality and embryonic stem cells generate more proliferative and vascularized teratomas ²¹
	−/−	Embryonic lethality due to (cardio)vascular defects and impaired tumor angiogenesis ^{26,27,81–83}
<i>HIF-2α</i>	+/-	Viable and fertile, protection against hypoxia-induced pulmonary vasculature remodeling, ⁸⁴ and reduced retinal neovascularization after oxygen-induced retinopathy ⁸⁵
	Endothelial inactivation	Defects in vessel integrity and tumor neovascularization ⁴¹
	−/− (Postnatal)	Anemia ⁸⁶
	−/− (Prenatal)	Background-dependent phenotype: embryonic lethality due to vascular defects and vascular disorganization in yolk sac and embryo (CD1 and ICR/129Sv outbred) ^{35,87} ; minimal viability due to multiple organ pathological features, metabolic abnormalities, and hematopoietic defects; defective catecholamine metabolism and bradycardia (C57/Black6 and 129 interbred) ^{37–39} ; defects in lung maturation; and acute respiratory distress (129/Sv x Swiss) ³⁶
<i>HIF-1α</i> ;; <i>HIF-2α</i>	−/−; −/−	Embryonic lethality due to defective placental vascularization ⁸⁸
<i>HIFdn</i>	Endothelial dominant negative	Embryonic lethality due to cardiovascular defects and an immature and disorganized vascular network ⁸⁹
<i>HIF-3α</i> (IPAS splice variant)	Oligo antisense in the cornea	Induced angiogenesis ⁹⁰
<i>HIF-3α</i> (NEPAS splice variant)	−/−	Enlargement of the right ventricle and impairment of lung remodeling ⁹¹
<i>HIF-1β</i>	Endothelial-specific inactivation	Partial embryonic lethality due to hepatic vascular defects and impaired energy homeostasis; and liver lesions, impaired lipid serum homeostasis, and hyperactive metabolism in adulthood ⁹²
	−/−	Embryonic lethality, insufficient angiogenesis and bronchial arch development, defective placental vascularization and vascular development, and altered placental cell fate determination ^{93–96}

ICR indicates ●●●; IPAS, ●●●; NEPAS, ●●●.

vated Notch1 and its target genes *Dtx1* and *Nrarp*.²¹ FIH-1 hydroxylates ankyrin repeat domains in Notch receptors, thereby decreasing their activity and the ability of FIH to hydroxylate HIFs.^{22,23}

Several genetic studies suggest a role for HIF-1α in vessel sprouting. Loss of HIF-1α in ECs attenuates the expression of hypoxia-inducible angiogenic genes and reduces EC proliferation under hypoxic conditions.^{24,25} EC-specific HIF-1α-deficient mice do not exhibit an overt phenotype in baseline conditions but show reduced tumor growth and vascularization because of attenuated endothelial expression of VEGF and VEGFR2. HIF-2α levels are not upregulated by the

absence of HIF-1α, suggesting that HIF-2α differentially regulates EC physiological features in vivo.²⁵ This is in line with previous reports^{26,27} that embryos lacking HIF-1α display vascular defects that are not rescued by HIF-2α. Conversely, HIF-1α stimulates neovascularization and improves functional recovery of ischemic tissues in adults.²⁸ Thus, although HIF-1α controls vessel sprouting, HIF-2α seems to mediate the maintenance of the newly formed sprout.

Oxygen Sensing and Endothelial Quiescence

Once a functional branch is established and is ready to fuse, the explorative behavior of the tip cell and the proliferative

nature of the stalk cell need to be suppressed. Recent studies report the involvement of HIF/PHD oxygen sensors in vessel stabilization. Angiopoietin-1 induces normalization of immature vessels, a process associated with reduced PHD2 expression.²⁹ Via upregulation of HIF-2 α , haploinsufficiency of PHD2 also induces tumor vessel normalization.³⁰ PHD2 haploinsufficiency in tumor ECs does not affect vessel density, area, or lumen size; however, it “normalizes” the endothelial lining, barrier, and stability. This does not affect primary tumor growth. It does improve tumor oxygenation and establish a tight impenetrable endothelial barrier, which prevents a metastatic switch and impedes intravasation of tumor cells and metastasis, overall resulting in prolonged survival of tumor-bearing mice. Gene profiling revealed a lower induction of HIF-1 α -dependent metastatic genes and an upregulation of the junctional protein VE-cadherin and the VEGF-trap soluble VEGFR1.³⁰ VE-cadherin has induced a normalized endothelial phenotype indirectly by inhibiting VEGF-driven proliferation and survival.^{31,32} It also inhibits proliferation of ECs in cooperation with soluble VEGFR1 to induce EC quiescence.³³ Consistently, its expression is specifically upregulated by HIF-2 α but not by HIF-1 α .³⁴

Genetic HIF-2 α -deficient studies show variable phenotypes depending on the genetic background. In a CD1 strain, loss of HIF-2 α causes improper remodeling of nascent vessels into larger conduits,³⁵ whereas in other backgrounds, loss of HIF-2 α causes abnormal organ development or homeostasis without apparent vascular deficits.^{36–39} Silencing HIF-2 α by approximately 50% in host mice leads to the formation of an aberrant vascular network in tumors through reduced expression of Ephrin-A1 in ECs. This is not because of reduced sprouting vessels or hypoxic EC proliferation; rather, it is because of impaired remodeling of the microvasculature.⁴⁰ Consistently, HIF-2 α deficiency in ECs results in increased vessel permeability and defective tumor neovascularization by decreasing the expression of genes involved in adhesion and extracellular matrix formation and angiogenesis.⁴¹ Notably, loss of HIF-2 α in ECs reduces metastasis (possibly because primary tumor growth was also reduced),⁴¹ whereas HIF-2 α upregulation in PHD2-haploinsufficient mice also reduces metastasis (via the previously mentioned mechanism of tumor vessel normalization).³⁰ In contrast, PHD2 inactivation in tumor cells accelerates tumor growth via HIF-independent NF- κ B pathway signaling.⁴² Also, complete inactivation of PHD2 in mice after birth stimulates the outgrowth of supernumerary vessels in healthy organs,⁴³ indicating that the effects of PHD2 in the vasculature are dose dependent.

Bone Marrow–Derived Progenitors, Oxygen Sensing, and Angiogenesis

Recruitment of a heterogeneous population of bone marrow-derived cells (BMDCs), containing progenitors of the endothelial, pericyte, and hematopoietic lineage, may also contribute to vessel formation and maturation.⁴⁴ Some of the signals that regulate the mobilization and homing of BMDCs to hypoxic areas have been characterized. VEGFR1 and its ligands (ie, VEGF and PlGF) induce BMDC mobilization and recruitment to ischemic regions^{45–47} in response to HIF-1 α stabilization.^{48,49} By binding CXCR4-positive BMDCs, stro-

mal cell–derived factor-1 (SDF-1/CXCL12) promotes recruitment and retention of endothelial progenitor cells in ischemic tissues. HIF-1 α upregulates CXCR4 receptor expression and enhances SDF-1 expression in ECs, thereby promoting adhesion, migration, and retention of CXCR4⁺ BMDCs.^{13,50} The inhibition of SDF-1 in ischemic tissue or of CXCR4 in circulating cells prevents endothelial progenitor cell recruitment to sites of injury.⁵¹ HIF-1 α -dependent induction of SDF-1 was documented in various angiogenesis models.^{52,53} Interestingly, hypoxic BM regions also exhibit increased SDF-1 expression and endothelial progenitor cell retention, suggesting that tissue hypoxia may constitute a more general mechanism governing stem cell retention.⁵¹

HIF-1 α -dependent induction of SDF-1 is also capable of recruiting bone marrow–derived CD45⁺ cells of the monocytic lineage.⁵⁴ Monocyte-derived cells secrete matrix metalloproteases, which contribute to angiogenesis by degrading basement membranes and extracellular matrix components, thereby enhancing the release of proangiogenic factors from the extracellular matrix.^{55–58} In addition, macrophages release the peptide PR39, which inhibits HIF-1 α degradation, amplifying the hypoxic response.⁵⁹ Hypoxic induction of matrix metalloproteases contributed to cancer cell metastasis by degrading extracellular matrix components.^{60–66} Development of hypoxia during antiangiogenic treatment may trigger recruitment of BMDCs, which can contribute to revascularization after treatment cessation. Interestingly, recent evidence links reduced tumor vascularization and growth during chemotherapy with a reduction in BMDC mobilization. Anthracyclines inhibit HIF-1 α activity by disrupting its binding to DNA, thereby decreasing levels of HIF-1 α target genes (eg, VEGF, SDF-1 α , and stem cell factor).⁶⁷

Mural Cells, Oxygen Sensing, and Vessel Maturation

Mural cells (ie, pericytes in capillaries and vascular smooth muscle cells in larger blood vessels) provide vessel stabilization and regulate vessel perfusion.⁶⁸ In PHD2-haploinsufficient mice, the reduction of PHD2 levels in ECs promotes pericyte coverage of tumor ECs via HIF-2 α -dependent mechanisms, thereby rendering the vessels more mature, tight, and stable.³⁰ In another experimental setting, inhibition of HIF-1 α reduces mural cell association with tumor ECs.⁶⁹ Furthermore, noise-induced mechanical and metabolic stress causes local hypoxia in the cochlear microvasculature, which promotes the stabilization of HIF-1 α . The subsequent upregulation of VEGF in both pericytes and ECs results in exaggerated pericyte coverage and abnormal vessel morphological features, yielding leaky and dysfunctional vessels.⁷⁰ Another link between oxygen sensing and vessel maturation may be platelet-derived growth factor receptor- β signaling, which controls mural cell recruitment and can be suppressed by VEGF-induced activation of VEGFR2.⁷¹

Conclusions

An important trigger activating the formation of a new blood vessel is the lack of oxygen in surrounding tissues. Therefore, it is not surprising that ECs and other cells composing the mature vessel are able to sense oxygen. Although previous

research has mainly focused on the role of VEGF, evidence is emerging that oxygen levels inside the vessel wall itself determine the fate of the developing sprout. Therefore, exploring the role of HIF- α , a master modulator of the hypoxic response, and its regulators (ie, PHD oxygen sensors) is crucial to improving our understanding of angiogenesis. Inside the endothelium, HIF-1 α is presumed to control sprouting, whereas HIF-2 α , in a dose-dependent way, ensures vessel quiescence and stability. In mural cells, a similar dose dependency also seems to determine the HIF-1 α -mediated response to hypoxia; however, it is still unclear whether HIF-2 α is also involved in conferring adequate pericyte or smooth muscle cell function. In addition, HIF-1 α has triggered the recruitment of progenitor cells that could also contribute to vessel stabilization.

The knowledge that the oxygen machinery influences all players conferring a mature functional vessel has broad clinical implications. Increasing tumor hypoxia through anti-angiogenic therapy induces the PHD/HIF oxygen-sensing machinery that is likely to reestablish the vasculature as soon as therapy ceases. Targeting the HIF transcriptional system in selective cell types may provide an elegant way to overcome this survival mechanism. Furthermore, treatment of diseases characterized by insufficient angiogenesis could also benefit from a cell-specific modulation of the HIF/PHD system. Therefore, more research is required to characterize the molecular pathways initiated by the oxygen sensors.

Sources of Funding

This study was supported by the Institute for the Promotion of Innovation by Science and Technology in Flanders (Dr Coulon); Leducq Network of Excellence in Atherothrombosis (Dr Roncal); the Fund for Scientific Research in Flanders (Dr De Bock); long-term structural funding (Methusalem funding by the Flemish government) (Dr Carmeliet); interuniversity attraction pole (grant P60/30 from the Belgian government) (Dr Carmeliet); the Fund for Scientific Research in Flanders G.0692.09 (Flemish government) (Dr Carmeliet); and research grant GAO 2006/11 from the Belgian Foundation Against Cancer (Dr Carmeliet).

Disclosures

The authors disclose a possible conflict of interest (patent WO2010084134 [A1]).

References

- Kaelin WG Jr, Ratcliffe PJ. Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. *Mol Cell*. 2008;30:393–402.
- Semenza GL. Targeting HIF-1 for cancer therapy. *Nat Rev Cancer*. 2003;3:721–732.
- Mahon PC, Hirota K, Semenza GL. FIH-1: a novel protein that interacts with HIF-1 α and VHL to mediate repression of HIF-1 transcriptional activity. *Genes Dev*. 2001;15:2675–2686.
- Fong GH. Regulation of angiogenesis by oxygen sensing mechanisms. *J Mol Med*. 2009.
- Takeda K, Ho VC, Takeda H, Duan LJ, Nagy A, Fong GH. Placental but not heart defects are associated with elevated hypoxia-inducible factor α levels in mice lacking prolyl hydroxylase domain protein 2. *Mol Cell Biol*. 2006;26:8336–8346.
- Berra E, Benizri E, Ginouves A, Volmat V, Roux D, Pouyssegur J. HIF prolyl-hydroxylase 2 is the key oxygen sensor setting low steady-state levels of HIF-1 α in normoxia. *EMBO J*. 2003;22:4082–4090.
- Yee Koh M, Spivak-Kroizman TR, Powis G. HIF-1 regulation: not so easy come, easy go. *Trends Biochem Sci*. 2008;33:526–534.
- Gerhardt H, Golding M, Fruttiger M, Ruhrberg C, Lundkvist A, Abramsson A, Jeltsch M, Mitchell C, Alitalo K, Shima D, Betsholtz C. VEGF guides angiogenic sprouting utilizing endothelial tip cell filopodia. *J Cell Biol*. 2003;161:1163–1177.
- Adams RH, Alitalo K. Molecular regulation of angiogenesis and lymphangiogenesis. *Nat Rev Mol Cell Biol*. 2007;8:464–478.
- Lu X, Le Noble F, Yuan L, Jiang Q, De Lafarge B, Sugiyama D, Breant C, Claes F, De Smet F, Thomas JL, Autiero M, Carmeliet P, Tessier-Lavigne M, Eichmann A. The netrin receptor UNC5B mediates guidance events controlling morphogenesis of the vascular system. *Nature*. 2004;432:179–186.
- Weinstein BM. Vessels and nerves: marching to the same tune. *Cell*. 2005;120:299–302.
- Unoki N, Murakami T, Nishijima K, Ogino K, van Rooijen N, Yoshimura N. SDF-1/CXCR4 contributes to the activation of tip cells and microglia in retinal angiogenesis. *Invest Ophthalmol Vis Sci*.
- Strasser GA, Kaminker JS, Tessier-Lavigne M. Microarray analysis of retinal endothelial tip cells identifies CXCR4 as a mediator of tip cell morphology and branching. *Blood*.
- Wang Y, Nakayama M, Pitulescu ME, Schmidt TS, Bochenek ML, Sakakibara A, Adams S, Davy A, Deutsch U, Luthi U, Barberis A, Benjamin LE, Makinen T, Nobes CD, Adams RH. Ephrin-B2 controls VEGF-induced angiogenesis and lymphangiogenesis. *Nature*. 465:483–486.
- Sawamiphak S, Seidel S, Essmann CL, Wilkinson GA, Pitulescu ME, Acker T, Acker-Palmer A. Ephrin-B2 regulates VEGFR2 function in developmental and tumour angiogenesis. *Nature*. 465:487–491.
- De Smet F, Segura I, De Bock K, Hohensinner PJ, Carmeliet P. Mechanisms of vessel branching: filopodia on endothelial tip cells lead the way. *Arterioscler Thromb Vasc Biol*. 2009;29:639–649.
- Hellstrom M, Phng LK, Hofmann JJ, Wallgard E, Coultas L, Lindblom P, Alva J, Nilsson AK, Karlsson L, Gaiano N, Yoon K, Rossant J, Iruela-Arispe ML, Kalen M, Gerhardt H, Betsholtz C. Dll4 signalling through Notch1 regulates formation of tip cells during angiogenesis. *Nature*. 2007;445:776–780.
- Gustafsson MV, Zheng X, Pereira T, Gradin K, Jin S, Lundkvist J, Ruas JL, Poellinger L, Lendahl U, Bondesson M. Hypoxia requires notch signaling to maintain the undifferentiated cell state. *Dev Cell*. 2005;9:617–628.
- Diez H, Fischer A, Winkler A, Hu CJ, Hatzopoulos AK, Breier G, Gessler M. Hypoxia-mediated activation of Dll4-Notch-Hey2 signaling in endothelial progenitor cells and adoption of arterial cell fate. *Exp Cell Res*. 2007;313:1–9.
- Patel NS, Li J-L, Generali D, Poulsom R, Cranston DW, Harris AL. Up-regulation of delta-like 4 ligand in human tumor vasculature and the role of basal expression in endothelial cell function. *Cancer Res*. 2005;65:8690–8697.
- Bertout JA, Patel SA, Fryer BH, Durham AC, Covellos KL, Olive KP, Goldschmidt MH, Simon MC. Heterozygosity for hypoxia inducible factor 1 α decreases the incidence of thymic lymphomas in a p53 mutant mouse model. *Cancer Res*. 2009;69:3213–3220.
- Coleman ML, McDonough MA, Hewitson KS, Coles C, Mecinovic J, Edelmann M, Cook KM, Cockman ME, Lancaster DE, Kessler BM, Oldham NJ, Ratcliffe PJ, Schofield CJ. Asparaginyl hydroxylation of the Notch ankyrin repeat domain by factor inhibiting hypoxia-inducible factor. *J Biol Chem*. 2007;282:24027–24038.
- Zheng X, Linke S, Dias JM, Gradin K, Wallis TP, Hamilton BR, Gustafsson M, Ruas JL, Wilkins S, Bilton RL, Brismar K, Whitelaw ML, Pereira T, Gorman JJ, Ericson J, Peet DJ, Lendahl U, Poellinger L. Interaction with factor inhibiting HIF-1 defines an additional mode of cross-coupling between the Notch and hypoxia signaling pathways. *Proc Natl Acad Sci U S A*. 2008;105:3368–3373.
- Sowter HM, Raval RR, Moore JW, Ratcliffe PJ, Harris AL. Predominant role of hypoxia-inducible transcription factor (Hif)-1 α versus Hif-2 α in regulation of the transcriptional response to hypoxia. *Cancer Res*. 2003;63:6130–6134.
- Tang N, Wang L, Esko J, Giordano FJ, Huang Y, Gerber HP, Ferrara N, Johnson RS. Loss of HIF-1 α in endothelial cells disrupts a hypoxia-driven VEGF autocrine loop necessary for tumorigenesis. *Cancer Cell*. 2004;6:485–495.
- Carmeliet P, Dor Y, Herbert JM, Fukumura D, Brusselmans K, Dewerchin M, Neeman M, Bono F, Abramovitch R, Maxwell P, Koch CJ, Ratcliffe P, Moons L, Jain RK, Collen D, Keshert E. Role of HIF-1 α in hypoxia-mediated apoptosis, cell proliferation and tumour angiogenesis. *Nature*. 1998;394:485–490.
- Iyer NV, Kotch LE, Agani F, Leung SW, Laughner E, Wenger RH, Gassmann M, Gearhart JD, Lawler AM, Yu AY, Semenza GL. Cellular

- and developmental control of O₂ homeostasis by hypoxia-inducible factor 1 alpha. *Genes Dev.* 1998;12:149–162.
28. Vincent KA, Shyu KG, Luo Y, Magner M, Tio RA, Jiang C, Goldberg MA, Akita GY, Gregory RJ, Isner JM. Angiogenesis is induced in a rabbit model of hindlimb ischemia by naked DNA encoding an HIF-1alpha/VP16 hybrid transcription factor. *Circulation.* 2000;102:2255–2261.
 29. Chen JX, Stinnett A. Ang-1 gene therapy inhibits hypoxia-inducible factor-1alpha (HIF-1alpha)-prolyl-4-hydroxylase-2, stabilizes HIF-1alpha expression, and normalizes immature vasculature in db/db mice. *Diabetes.* 2008;57:3335–3343.
 30. Mazzone M, Dettori D, Leite de Oliveira R, Loges S, Schmidt T, Jonckx B, Tian Y-M, Lanahan AA, Pollard P, Ruiz de Almodovar C, De Smet F, Vinckier S, Aragonès JN, Debackere K, Luttun A, Wyns S, Jordan B, Pisacane A, Gallez B, Lampugnani MG, Dejana E, Simons M, Ratcliffe P, Maxwell P, Carmeliet P. Heterozygous deficiency of PHD2 restores tumor oxygenation and inhibits metastasis via endothelial normalization. *Cell.* 2009;136:839–851.
 31. Carmeliet P, Lampugnani MG, Moons L, Breviaro F, Compennolle V, Bono F, Balconi G, Spagnuolo R, Oosthuysen B, Dewerchin M, Zanetti A, Angelillo A, Mattot V, Nuyens D, Lutgens E, Clotman F, de Ruiter MC, Gittenberger-de Groot A, Poelmann R, Lupu F, Herbert JM, Collen D, Dejana E. Targeted deficiency or cytosolic truncation of the VE-cadherin gene in mice impairs VEGF-mediated endothelial survival and angiogenesis. *Cell.* 1999;98:147–157.
 32. Nakamura Y, Patrushev N, Inomata H, Mehta D, Urao N, Kim HW, Razvi M, Kini V, Mahadev K, Goldstein BJ, McKinney R, Fukai T, Ushio-Fukai M. Role of protein tyrosine phosphatase 1B in vascular endothelial growth factor signaling and cell-cell adhesions in endothelial cells. *Circ Res.* 2008;102:1182–1191.
 33. Kearney JB, Kappas NC, Ellerstrom C, DiPaola FW, Bautch VL. The VEGF receptor flt-1 (VEGFR-1) is a positive modulator of vascular sprout formation and branching morphogenesis. *Blood.* 2004;103:4527–4535.
 34. Le Bras A, Lionneton F, Mattot V, Lelievre E, Caetano B, Spruyt N, Soncin F. HIF-2alpha specifically activates the VE-cadherin promoter independently of hypoxia and in synergy with Ets-1 through two essential ETS-binding sites. *Oncogene.* 2007;26:7480–7489.
 35. Duan LJ, Zhang-Benoit Y, Fong GH. Endothelium-intrinsic requirement for Hif-2alpha during vascular development. *Circulation.* 2005;111:2227–2232.
 36. Compennolle V, Brusselmans K, Acker T, Hoet P, Tjwa M, Beck H, Plaisance S, Dor Y, Keshet E, Lupu F, Nemery B, Dewerchin M, Van Veldhoven P, Plate K, Moons L, Collen D, Carmeliet P. Loss of HIF-2alpha and inhibition of VEGF impair fetal lung maturation, whereas treatment with VEGF prevents fatal respiratory distress in premature mice. *Nat Med.* 2002;8:702–710.
 37. Scortegagna M, Ding K, Oktay Y, Gaur A, Thurmond F, Yan LJ, Marck BT, Matsumoto AM, Shelton JM, Richardson JA, Bennett MJ, Garcia JA. Multiple organ pathology, metabolic abnormalities and impaired homeostasis of reactive oxygen species in Epas1^{-/-} mice. *Nat Genet.* 2003;35:331–340.
 38. Scortegagna M, Morris MA, Oktay Y, Bennett M, Garcia JA. The HIF family member EPAS1/HIF-2alpha is required for normal hematopoiesis in mice. *Blood.* 2003;102:1634–1640.
 39. Tian H, Hammer RE, Matsumoto AM, Russell DW, McKnight SL. The hypoxia-responsive transcription factor EPAS1 is essential for catecholamine homeostasis and protection against heart failure during embryonic development. *Genes Dev.* 1998;12:3320–3324.
 40. Yamashita T, Ohneda K, Nagano M, Miyoshi C, Kaneko N, Miwa Y, Yamamoto M, Ohneda O, Fujii-Kuriyama Y. HIF-2alpha in endothelial cells regulates tumor neovascularization through activation of ephrin A1. *J Biol Chem.* 2008;18926–18936.
 41. Skuli N, Liu L, Runge A, Wang T, Yuan L, Patel S, Iruela-Arispe L, Simon MC, Keith B. Endothelial deletion of hypoxia inducible factor-2alpha (HIF-2{alpha}) alters vascular function and tumor angiogenesis. *Blood.* 2009.
 42. Chan DA, Kawahara TL, Sutphin PD, Chang HY, Chi JT, Giaccia AJ. Tumor vasculature is regulated by PHD2-mediated angiogenesis and bone marrow-derived cell recruitment. *Cancer Cell.* 2009;15:527–538.
 43. Takeda K, Aguila HL, Parikh NS, Li X, Lamothe K, Duan L-J, Takeda H, Lee FS, Fong G-H. Regulation of adult erythropoiesis by prolyl hydroxylase domain proteins. *Blood.* 2008;111:3229–3235.
 44. Ahn GO, Brown JM. Role of endothelial progenitors and other bone marrow-derived cells in the development of the tumor vasculature. *Angiogenesis.* 2009;12:159–164.
 45. Carmeliet P, Moons L, Luttun A, Vincenti V, Compennolle V, De Mol M, Wu Y, Bono F, Devy L, Beck H, Scholz D, Acker T, DiPalma T, Dewerchin M, Noel A, Stalmans I, Barra A, Blacher S, Vandendriessche T, Ponten A, Eriksson U, Plate KH, Foidart JM, Schaper W, Charnock-Jones DS, Hicklin DJ, Herbert JM, Collen D, Persico MG. Synergism between vascular endothelial growth factor and placental growth factor contributes to angiogenesis and plasma extravasation in pathological conditions. *Nat Med.* 2001;7:575–583.
 46. Luttun A, Tjwa M, Moons L, Wu Y, Angelillo-Scherrer A, Liao F, Nagy JA, Hooper A, Priller J, De Klerck B, Compennolle V, Daci E, Bohlen P, Dewerchin M, Herbert JM, Fava R, Matthys P, Carmeliet G, Collen D, Dvorak HF, Hicklin DJ, Carmeliet P. Revascularization of ischemic tissues by PIGF treatment, and inhibition of tumor angiogenesis, arthritis and atherosclerosis by anti-Flt1. *Nat Med.* 2002;8:831–840.
 47. Hattori K, Heissig B, Wu Y, Dias S, Tejada R, Ferris B, Hicklin DJ, Zhu Z, Bohlen P, Witte L, Hendriks J, Hackett NR, Crystal RG, Moore MA, Werb Z, Lyden D, Rafii S. Placental growth factor reconstitutes hematopoiesis by recruiting VEGFR1(+) stem cells from bone-marrow microenvironment. *Nat Med.* 2002;8:841–849.
 48. Liu L, Marti GP, Wei X, Zhang X, Zhang H, Liu YV, Nastai M, Semenza GL, Harmon JW. Age-dependent impairment of HIF-1alpha expression in diabetic mice: correction with electroporation-facilitated gene therapy increases wound healing, angiogenesis, and circulating angiogenic cells. *J Cell Physiol.* 2008;217:319–327.
 49. Okuyama H, Krishnamachary B, Zhou YF, Nagasawa H, Bosch-Marce M, Semenza GL. Expression of vascular endothelial growth factor receptor 1 in bone marrow-derived mesenchymal cells is dependent on hypoxia-inducible factor 1. *J Biol Chem.* 2006;281:15554–15563.
 50. Staller P, Sulitkova J, Lisztwan J, Moch H, Oakeley EJ, Krek W. Chemokine receptor CXCR4 downregulated by von Hippel-Lindau tumour suppressor pVHL. *Nature.* 2003;425:307–311.
 51. Ceradini DJ, Kulkarni AR, Callaghan MJ, Tepper OM, Bastidas N, Kleinman ME, Capla JM, Galiano RD, Levine JP, Gurtner GC. Progenitor cell trafficking is regulated by hypoxic gradients through HIF-1 induction of SDF-1. *Nat Med.* 2004;10:858–864.
 52. Ritter MR, Banin E, Moreno SK, Aguilar E, Dorrell MI, Friedlander M. Myeloid progenitors differentiate into microglia and promote vascular repair in a model of ischemic retinopathy. *J Clin Invest.* 2006;116:3266–3276.
 53. Fadini GP, Sartore S, Schiavon M, Albiero M, Baesso I, Cabrelle A, Agostini C, Avogaro A. Diabetes impairs progenitor cell mobilisation after hindlimb ischaemia-reperfusion injury in rats. *Diabetologia.* 2006;49:3075–3084.
 54. Du R, Lu KV, Petritsch C, Liu P, Ganss R, Passegue E, Song H, Vandenberg S, Johnson RS, Werb Z, Bergers G. HIF1alpha induces the recruitment of bone marrow-derived vascular modulatory cells to regulate tumor angiogenesis and invasion. *Cancer Cell.* 2008;13:206–220.
 55. Jodele S, Chantre CF, Blavier L, Lutzko C, Crooks GM, Shimada H, Coussens LM, Declercq YA. The contribution of bone marrow-derived cells to the tumor vasculature in neuroblastoma is matrix metalloproteinase-9 dependent. *Cancer Res.* 2005;65:3200–3208.
 56. Nozawa H, Chiu C, Hanahan D. Infiltrating neutrophils mediate the initial angiogenic switch in a mouse model of multistage carcinogenesis. *Proc Natl Acad Sci U S A.* 2006;103:12493–12498.
 57. Page-McCaw A, Ewald AJ, Werb Z. Matrix metalloproteinases and the regulation of tissue remodelling. *Nat Rev Mol Cell Biol.* 2007;8:221–233.
 58. Yang L, DeBusk LM, Fukuda K, Fingleton B, Green-Jarvis B, Shyr Y, Matrisian LM, Carbone DP, Lin PC. Expansion of myeloid immune suppressor Gr+CD11b+ cells in tumor-bearing host directly promotes tumor angiogenesis. *Cancer Cell.* 2004;6:409–421.
 59. Li J, Post M, Volk R, Gao Y, Li M, Metais C, Sato K, Tsai J, Aird W, Rosenberg RD, Hampton TG, Sellke F, Carmeliet P, Simons M. PR39, a peptide regulator of angiogenesis. *Nat Med.* 2000;6:49–55.
 60. Wu H, Liang X, Fang Y, Qin X, Zhang Y, Liu J. Resveratrol inhibits hypoxia-induced metastasis potential enhancement by restricting hypoxia-induced factor-1 alpha expression in colon carcinoma cells. *Biomed Pharmacother.* 2008;62:613–621.
 61. Miyazaki Y, Hara A, Kato K, Oyama T, Yamada Y, Mori H, Shibata T. The effect of hypoxic microenvironment on matrix metalloproteinase expression in xenografts of human oral squamous cell carcinoma. *Int J Oncol.* 2008;32:145–151.
 62. Fujiwara S, Nakagawa K, Harada H, Nagato S, Furukawa K, Teraoka M, Seno T, Oka K, Iwata S, Ohnishi T. Silencing hypoxia-inducible factor-1alpha inhibits cell migration and invasion under hypoxic environment in malignant gliomas. *Int J Oncol.* 2007;30:793–802.

63. Spinella F, Rosano L, Di Castro V, Decandia S, Nicotra MR, Natali PG, Bagnato A. Endothelin-1 and endothelin-3 promote invasive behavior via hypoxia-inducible factor-1 α in human melanoma cells. *Cancer Res*. 2007;67:1725–1734.
64. Lungu GF, Li ML, Xie X, Wang LV, Stoica G. In vivo imaging and characterization of hypoxia-induced neovascularization and tumor invasion. *Int J Oncol*. 2007;30:45–54.
65. Petrella BL, Brinckerhoff CE. Tumor cell invasion of von Hippel Lindau renal cell carcinoma cells is mediated by membrane type-1 matrix metalloproteinase. *Mol Cancer*. 2006;5:66.
66. Luo Y, He DL, Ning L, Shen SL, Li L, Li X, Zhou HE, Chung LW. Over-expression of hypoxia-inducible factor-1 α increases the invasive potency of LNCaP cells in vitro. *BJU Int*. 2006;98:1315–1319.
67. Lee K, Qian DZ, Rey S, Wei H, Liu JO, Semenza GL. Anthracycline chemotherapy inhibits HIF-1 transcriptional activity and tumor-induced mobilization of circulating angiogenic cells. *Proc Natl Acad Sci U S A*. 2009;106:2353–2358.
68. Carmeliet P. Angiogenesis in life, disease and medicine. *Nature*. 2005;438:932–936.
69. Stoeltz O, McCarty MF, Wey JS, Fan F, Liu W, Belcheva A, Bucana CD, Semenza GL, Ellis LM. Role of hypoxia-inducible factor 1 α in gastric cancer cell growth, angiogenesis, and vessel maturation. *J Natl Cancer Inst*. 2004;96:946–956.
70. Shi X. Cochlear pericyte responses to acoustic trauma and the involvement of hypoxia-inducible factor-1 α and vascular endothelial growth factor. *Am J Pathol*. 2009.
71. Greenberg JJ, Shields DJ, Barillas SG, Acevedo LM, Murphy E, Huang J, Scheppke L, Stockmann C, Johnson RS, Angle N, Cheresch DA. A role for VEGF as a negative regulator of pericyte function and vessel maturation. *Nature*. 2008;456:809–813.
72. Aragonés J, Schneider M, Van Geyte K, Fraisl P, Dresselaers T, Mazzone M, Dirx R, Zaccagna S, Lemieux H, Jeoung NH, Lambrechts D, Bishop T, Lafuste P, Diez-Juan A, Harten SK, Van Noten P, De Bock K, Willam C, Tjwa M, Grosfeld A, Navet R, Moons L, Vandendriessche T, Deroose C, Wijeyekoon B, Nuyts J, Jordan B, Silasi-Mansat R, Lupu F, Dewerchin M, Pugh C, Salmon P, Mortelmans L, Gallez B, Goris F, Buyse J, Sluse F, Harris RA, Gnaiger E, Hespel P, Van Hecke P, Schuit F, Van Veldhoven P, Ratcliffe P, Baes M, Maxwell P, Carmeliet P. Deficiency or inhibition of oxygen sensor PHD1 induces hypoxia tolerance by reprogramming basal metabolism. *Nat Genet*. 2008;40:170–180.
73. Schneider M, Van Geyte K, Fraisl P, Kiss J, Aragonés J, Mazzone M, Mairbaurl H, De Bock K, Jeoung NH, Mollenhauer M, Georgiadou M, Bishop T, Roncal C, Sutherland A, Jordan B, Gallez B, Weitz J, Harris RA, Maxwell P, Baes M, Ratcliffe P, Carmeliet P. Loss or silencing of the PHD1 prolyl hydroxylase protects livers of mice against ischemia/reperfusion injury. *Gastroenterology*. 138:1143–1154, e1141–e1142.
74. Zhang Q, Gu J, Li L, Liu J, Luo B, Cheung HW, Boehm JS, Ni M, Geisen C, Root DE, Polyak K, Brown M, Richardson AL, Hahn WC, Kaelin WG Jr, Bommi-Reddy A. Control of cyclin D1 and breast tumorigenesis by the EglN2 prolyl hydroxylase. *Cancer Cell*. 2009;16:413–424.
75. Takeda K, Cowan A, Fong GH. Essential role for prolyl hydroxylase domain protein 2 in oxygen homeostasis of the adult vascular system. *Circulation*. 2007;116:774–781.
76. Minamishima YA, Moslehi J, Bardeesy N, Cullen D, Bronson RT, Kaelin WG Jr. Somatic inactivation of the PHD2 prolyl hydroxylase causes polycythemia and congestive heart failure. *Blood*. 2008;111:3236–3244.
77. Bishop T, Gallagher D, Pascual A, Lygate CA, de Bono JP, Nicholls LG, Ortega-Saenz P, Oster H, Wijeyekoon B, Sutherland AI, Grosfeld A, Aragonés J, Schneider M, van Geyte K, Teixeira D, Diez-Juan A, Lopez-Barneo J, Channon KM, Maxwell PH, Pugh CW, Davies AM, Carmeliet P, Ratcliffe PJ. Abnormal sympathoadrenal development and systemic hypotension in PHD3 $^{-/-}$ mice. *Mol Cell Biol*. 2008;28:3386–3400.
78. Minamishima YA, Moslehi J, Padera RF, Bronson RT, Liao R, Kaelin WG Jr. A feedback loop involving the PHD3 prolyl hydroxylase tunes the mammalian hypoxic response in vivo. *Mol Cell Biol*. 2009;29:5729–5741.
79. Zhang N, Fu Z, Linke S, Chicher J, Gorman JJ, Visk D, Haddad GG, Poellinger L, Peet DJ, Powell F, Johnson RS. The asparaginyl hydroxylase factor inhibiting HIF-1 α is an essential regulator of metabolism. *Cell Metab*. 11:364–378.
80. Yu AY, Shimoda LA, Iyer NV, Huso DL, Sun X, McWilliams R, Beaty T, Sham JS, Wiener CM, Sylvester JT, Semenza GL. Impaired physiological responses to chronic hypoxia in mice partially deficient for hypoxia-inducible factor 1 α . *J Clin Invest*. 1999;103:691–696.
81. Ryan HE, Lo J, Johnson RS. HIF-1 α is required for solid tumor formation and embryonic vascularization. *EMBO J*. 1998;17:3005–3015.
82. Kotch LE, Iyer NV, Laughner E, Semenza GL. Defective vascularization of HIF-1 α -null embryos is not associated with VEGF deficiency but with mesenchymal cell death. *Dev Biol*. 1999;209:254–267.
83. Compennolle V, Brusselmans K, Franco D, Moorman A, Dewerchin M, Collen D, Carmeliet P. Cardiac bifida, defective heart development and abnormal neural crest migration in embryos lacking hypoxia-inducible factor-1 α . *Cardiovasc Res*. 2003;60:569–579.
84. Brusselmans K, Compennolle V, Tjwa M, Wiesener MS, Maxwell PH, Collen D, Carmeliet P. Heterozygous deficiency of hypoxia-inducible factor-2 α protects mice against pulmonary hypertension and right ventricular dysfunction during prolonged hypoxia. *J Clin Invest*. 2003;111:1519–1527.
85. Dioum EM, Clarke SL, Ding K, Repa JJ, Garcia JA. HIF-2 α -haploinsufficient mice have blunted retinal neovascularization due to impaired expression of a proangiogenic gene battery. *Invest Ophthalmol Vis Sci*. 2008;49:2714–2720.
86. Gruber M, Hu CJ, Johnson RS, Brown EJ, Keith B, Simon MC. Acute postnatal ablation of Hif-2 α results in anemia. *Proc Natl Acad Sci U S A*. 2007;104:2301–2306.
87. Peng J, Zhang L, Drysdale L, Fong GH. The transcription factor EPAS-1/hypoxia-inducible factor 2 α plays an important role in vascular remodeling. *Proc Natl Acad Sci U S A*. 2000;97:8386–8391.
88. Cowden Dahl KD, Robertson SE, Weaver VM, Simon MC. Hypoxia-inducible factor regulates α 5 β 3 integrin cell surface expression. *Mol Biol Cell*. 2005;16:1901–1912.
89. Licht AH, Muller-Holtkamp F, Flamme I, Breier G. Inhibition of hypoxia-inducible factor activity in endothelial cells disrupts embryonic cardiovascular development. *Blood*. 2006;107:584–590.
90. Makino Y, Cao R, Svensson K, Bertilsson G, Asman M, Tanaka H, Cao Y, Berkenstam A, Poellinger L. Inhibitory PAS domain protein is a negative regulator of hypoxia-inducible gene expression. *Nature*. 2001;414:550–554.
91. Yamashita T, Ohneda O, Nagano M, Iemitsu M, Makino Y, Tanaka H, Miyauchi T, Goto K, Ohneda K, Fujii-Kuriyama Y, Poellinger L, Yamamoto M. Abnormal heart development and lung remodeling in mice lacking the hypoxia-inducible factor-related basic helix-loop-helix PAS protein NEPAS. *Mol Cell Biol*. 2008;28:1285–1297.
92. Yim SH, Shah Y, Tomita S, Morris HD, Gavrilova O, Lambert G, Ward JM, Gonzalez FJ. Disruption of the Arnt gene in endothelial cells causes hepatic vascular defects and partial embryonic lethality in mice. *Hepatology*. 2006;44:550–560.
93. Maltepe E, Schmidt JV, Baunoch D, Bradfield CA, Simon MC. Abnormal angiogenesis and responses to glucose and oxygen deprivation in mice lacking the protein ARNT. *Nature*. 1997;386:403–407.
94. Kozak KR, Abbott B, Hankinson O. ARNT-deficient mice and placental differentiation. *Dev Biol*. 1997;191:297–305.
95. Ramirez-Bergeron DL, Runge A, Adelman DM, Gohil M, Simon MC. HIF-dependent hematopoietic factors regulate the development of the embryonic vasculature. *Dev Cell*. 2006;11:81–92.
96. Adelman DM, Gertsenstein M, Nagy A, Simon MC, Maltepe E. Placental cell fates are regulated in vivo by HIF-mediated hypoxia responses. *Genes Dev*. 2000;14:3191–3203.